Fuel wood - Flow

General description

Not all forests can be managed equally: accessibility (infrastructure) and technical feasibility (due to topography) play a crucial role in sustainable forest management. Hence, this indicator reflects the timber removals for fuel wood production considering both the forest accessibility (based on the OSM layer) and topographical site condition (slope angle from DEM). Data for timber removals are taken from national forestry inventories. **Flowchart below** describes in detail the calculation procedure to derive the amount of fuelwood (m³/ha/year) produced per local administrative units (LAU2) of the Alpine Space.

The procedure consists of two main parts:

- Calculation of the accessible forest areas (Step 1 and 2): typically, commercial wood production occurs only on forests that are accessible and workable in terms of topographic conditions. Hence, in the following analysis only accessible and workable forest areas are included.
- <font 14px/inherit;;inherit;calculation of the amount of fuelwood produced in each LAU2 (Step 3 and 4): The statistical data about the amount of felling is disaggregated at LAU2 level using results from part 1.

Input data

GIS-Data

Forest roads: open street map layer (OSM)

Source: http://download.geofabrik.de/. The roads classified as "bridleway", "service", "track", "track_grade1", "track_grade2", "track_grade3", "track_grade4", "track_grade5", "unclassified" and "unknown" are used.

• Forest: High-resolution tree-cover-density layer

Source: http://land.copernicus.eu/. Pixels with a density between 30-100% are used; forest in agricultural and urban areas is removed.

Digital Elevation Model: Elevation raster dataset

Source: http://land.copernicus.eu,v1.0

• LAU2 layer: Boundaries of the local administrative units

Source: http://www.eurogeographics.org/products-and-services/euroboundarymap; version 10.

• **Region layer:**Boundaries of the areas for which the statistical data is available.Statistical data about the amount of felled wood is typically not

available at LAU level; hence, for every country we used the finest available resource. Because these datasets have different resolutions and names, we call them "region layers" in the following description. The original names/resolutions for the different countries are the following:

-Germany: Growing regions

Source: https://gdi.thuenen.de/wo/wgwb/

-Switzerland: Production regions*

-France: Nuts 2

Source: http://www.eurogeographics.org/products-and-services/euroboundarymap

-Italy: Nuts 2

Source: http://www.eurogeographics.org/products-and-services/euroboundarymap

-Liechtenstein: Nuts 2

Source: http://www.eurogeographics.org/products-and-services/euroboundarymap

-Austria: "Bezirke"

Source: http://data.opendataportal.at/dataset/geojson-daten-osterreich

-Slovenia: Forest regions (dataset provided by the Slovenian Forest Institute)

Statistical data<font

14px/inherit;;inherit>:

 Statistical data about roundwood production: The amount of wood that is extracted from the forest.

Source: Forest inventories of single countries

Percentage of roundwood transformed into fuelwood

Source:

http://ec.europa.eu/eurostat/statistics-explained/index.php/Wood products - production and trade

Variables:

Maximum access distance = 200m (after Clouet, N. and Berger, F., 2009)

Calculation processes

(1) Calculation of the slope angle of accessible forest: Forested areas within the maximum access distance from forest roads are extracted from the original forest layer and their slope is calculated (in %) The 200m serve as a threshold, all pixels that are further away from the street are not considered in the following calculation.

http://www.wikialps.eu/ Printed on 2025/07/12 08:58

2025/07/12 08.58 3/3 Fuel wood - Flow

From:
(2) Calculation of access distance: Based on Clouet & Berger (2009), the maxim which every pixel (extracted in Step 1) can be accessed, due to its topographic concareulased of the lower than the actual distance from the temporary who will pixel that the lower than the actual distance from the temporary will pixel that the lower than the actual distance from the temporary will pixel that the lower than the lower than

(3) apisaggregated to LAU2 level: The data about the production roundwood is disaggregated to LAU2 level via the share of accessible forest in every LAU2.