Section	Description
Resource Title	Filtration of surface water by ecosystem types – Flow/Supply
Resource Abstract	Filtration of surface water by terrestrial ecosystems is an important service that directly contributes to mitigate water pollution. This multi-faceted ES includes a variety of ecosystem types and pollutants. Among the different types of pollutants are nutrients, such as nitrogen or phosphorous, pesticides, heavy metals or organic matter. Within AlpES the outcomes of the InVEST NDR model (Nutrient Delivery Ratio) have been used to assess the nitrogen loads and the filtration capacities by different ecosystems. This model follows a mass balance approach, describing the movement of the mass of nutrient through space. Unlike more sophisticated nutrient models, NDR does not represent the details of the nutrient cycle but rather represents the long-term, steady-state flow of nutrients through empirical relationships.
Resource Type	Dataset
Resource locator	http://www.alpes- webgis.eu/?X=850359.92&Y=5947762.56&zoom=6⟨=en&focus=focus al pes&bgLayer=alpes.osm.stamentoner.60002&layers=alpes.alpinespace.40001. wms,alpes.essi.10072&catalogNodes=101000000,101000008&layers_opacity =1,0.7
Unique Resource Identifier	Z6WH-RYRJ-LQ56-CC3S
Resource Language	eng
Topic Category	Environment
Keyword value	Land use (INSPIRE Spatial Data Theme) Land cover (INSPIRE Spatial Data Theme) Filtration (GEMET Concepts Nitrogen (GEMET Concepts)
Originating	- title: GEMET - INSPIRE themes, version 1.0
controlled	- date:
vocabulary	-dateType: publication -date: 2008-06-01 - title: GEMET - Concepts, version 4.0.1 - date: -dateType: publication -date: 2017-06-28
Geographic bounding box	West = 1.986194 East = 18.622061
	North = 50.068114 South = 42.700501
Coordinate reference System	EPSG: 3035 (ETRS89, LAEA)
Temporal extent	2012

Date of publication	2018-07-20
Lineage	The flow/supply indicator represents the actual amount of nitrogen that is reduced within a single municipality. This is the capacity of nitrogen filtration, determined by the difference between the loaded nitrogen (N loads per land- use type, N atmospheric deposition and N biological fixation) and the nitrogen exported from the area of a municipality. Both these values are calculated using the InVEST NDR model. Units of measurements: kg ha <sup>-1</sup> y <sup>-1</sup>
Spatial resolution	100000
Specification	Commission Regulation (EU) No 1089/2010 of 23 November 2010
	implementing Directive 2007/2/EC of the European Parliament and of the
	Council as regards interoperability of spatial data sets and services, date of
	publication: 2010-12-08.
Degree	Null
<b>Conditions applying</b>	<u>CC BY-NC 4.0</u>
to access and use	
Limitations on public	No Limitation
access	
<b>Responsible party</b>	Eurac Research, Viale Druso 1, 39100 Bolzano, Italy
	Institute for Alpine Environment - <u>alpine.environment@eurac.edu</u>
Responsible party	Author
role	
Metadata point of	Eurac Research, Viale Druso 1, 39100 Bolzano, Italy
contact	Institute for Alpine Environment - <u>alpine.environment@eurac.edu</u>
Metadata date	2018-04-20
Metadata language	eng