Section	Description
Resource Title	Filtration of surface water by ecosystem types – Flow/Supply
Resource Abstract	Filtration of surface water by terrestrial ecosystems is an important service
	that directly contributes to mitigate water pollution. This multi-faceted ES
	includes a variety of ecosystem types and pollutants. Among the different
	types of pollutants are nutrients, such as nitrogen or phosphorous, pesticides,
	heavy metals or organic matter. Within AlpES the outcomes of the InVEST NDR
	model (Nutrient Delivery Ratio) have been used to assess the nitrogen loads
	and the filtration capacities by different ecosystems. This model follows a
	mass balance approach, describing the movement of the mass of nutrient
	through space. Unlike more sophisticated nutrient models, NDR does not
	represent the details of the nutrient cycle but rather represents the long-term,
	steady-state flow of nutrients through empirical relationships.
Resource Type	Dataset
Resource locator	http://www.alpes-
	webgis.eu/?X=850359.92&Y=5947762.56&zoom=6⟨=en&focus_al
	pes&bgLayer=alpes.osm.stamentoner.60002&layers=alpes.alpinespace.40001.
	wms,alpes.essi.10072&catalogNodes=101000000,101000008&layers_opacity
	=1,0.7
Unique Resource	Z6WH-RYRJ-LQ56-CC3S
Identifier	
Resource Language	eng
Topic Category	Environment
Keyword value	Land use (INSPIRE Spatial Data Theme)
	Land cover (INSPIRE Spatial Data Theme)
	Filtration (GEMET Concepts
	Nitrogen (GEMET Concepts)
Originating	- title: GEMET - INSPIRE themes, version 1.0
controlled	- date:
vocabulary	-dateType: publication
	-date: 2008-06-01
	- title: GEMET - Concepts, version 4.0.1
	- date:
	-dateType: publication -date: 2017-06-28
Coographic hounding	West = 1.986194
Geographic bounding box	West = 1.980194
DOX	East = 18.622061
	Last - 18.022001
	North = 50.068114
	101ul - 201000111
	South = 42.700501
Coordinate reference	EPSG: 3035 (ETRS89, LAEA)
System	
Temporal extent	From 2012-01-01T11:15:00 to 2012-12-31T11:15:00

Date of publication	2017-07-24 T11:15:00
Lineage	The flow/supply indicator represents the actual amount of nitrogen that is
	reduced within a single municipality. This is the capacity of nitrogen filtration,
	determined by the difference between the loaded nitrogen (N loads per land-
	use type, N atmospheric deposition and N biological fixation) and the nitrogen
	exported from the area of a municipality. Both these values are calculated
	using the InVEST NDR model.
Spatial resolution	100000
Specification	Commission Regulation (EU) No 1089/2010 of 23 November 2010
	implementing Directive 2007/2/EC of the European Parliament and of the
	Council as regards interoperability of spatial data sets and services, date of
	publication: 2010-12-08.
Degree	Null
Conditions applying	<u>CC BY-NC 4.0</u>
to access and use	
Limitations on public	No Limitation
access	
Responsible party	Eurac Research, Viale Druso 1, 39100 Bolzano, Italy
	Institute for Alpine Environment - <u>alpine.environment@eurac.edu</u>
Responsible party	Author
role	
Metadata point of	Eurac Research, Viale Druso 1, 39100 Bolzano, Italy
contact	Institute for Alpine Environment - <u>alpine.environment@eurac.edu</u>
Metadata date	2018-04-20
Metadata language	eng